

# Corvalen M®

## D-Ribose with Magnesium and Malate

### DESCRIPTION

Corvalen M® D-ribose, distributed exclusively to healthcare professionals by Douglas Laboratories, is a natural pentose sugar that is designed for the support of fatigue, energy production, and mitochondrial function<sup>†</sup>. Magnesium and malate are added to this formula to help in the utilization of energy and support symptoms of fatigue.


### FUNCTIONS

Corvalen M® contains pure D-ribose, a safe and clinically researched ingredient that supports the natural way our bodies produce adenosine triphosphate (ATP), the energy currency of the cell. Ribose is the vital structural backbone of critical cellular compounds called purines and pyrimidines. Our bodies must have an adequate supply of purines and pyrimidines to form major cellular constituents such as our genetic material (DNA and RNA), numerous cofactors, certain vitamins, and, importantly, adenosine triphosphate (ATP). Ribose is the starting point for the synthesis of these fundamental cellular compounds, and the availability of ribose determines the rate at which they can be made by our cells and tissues.

D-ribose is a structural component of DNA, RNA, ATP, GTP, flavins (FAD, riboflavin) and other important nucleotides found in all living cells. Ribose is formed naturally via the pentose phosphate pathway. This pathway is slow and rate-limited in cardiac and skeletal muscle due to an inherently low concentration (lack of expression) of the enzymes, glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase. The product of this pathway is ribose-5-phosphate, which in turn is converted to 5-phosphoribosyl-1-pyrophosphate (PRPP), the primary driver in the synthesis and salvage of purine nucleotides. No other compound can be used by the body for this metabolic purpose. Purine nucleotides (ATP and its precursors) lost due to ischemia, hypoxia, or genetic predisposition are replaced via the purine nucleotide pathway. This pathway is rate limited by the availability of ribose in tissue. Administration of exogenous ribose bypasses the rate-limiting steps in the pentose phosphate pathway, resulting in a significant acceleration of PRPP.



## Why Exogenous Ribose Speeds Energy Resynthesis



Corvalen M® D-ribose is highly soluble in both hot and cold solutions and tastes slightly sweet and tart. Corvalen M® D-ribose is non-GMO. D-ribose is rapidly and readily (~95%) absorbed with peak blood levels found within 30 – 45 minutes. Ribose not taken up by the cell is excreted unchanged in the urine. Corvalen® D-ribose is GRAS (generally recognized as safe), a determination that results only after considerable toxicology studies are performed by the FDA.

## INDICATIONS

Corvalen M® D-ribose is a useful dietary supplement for those patients that want to restore energy and support the symptoms of fatigue with the added benefit of magnesium and malate to help relieve occasionally sore muscles.

## **FORMULA (#57452)**

Serving Size 6.1 g (1 scoop or 2 tsp), serving per container 56

**D-ribose.....5 g**

**Magnesium Gluconate.....800 mg**

**(40 mg elemental magnesium)**

**Malate.....240 mg**

*No other ingredients.*

## **SUGGESTED USE**

**Usual dosage:** 1 scoop (2 tsp) serving twice daily, taken with meals. A third serving may be added with a mid-day meal or snack as needed.

**Alternative dosage:** 1 scoop 30 minutes before and just after exercise or physical activity.

Corvalen granular powder may be dissolved in 2 oz. or more of juice or liquid or sprinkled over other foods of choice. Do not mix with milk or carbonated beverages.

## **SIDE EFFECTS**

No adverse effects have been reported unless taking doses greater than 10 grams of ribose at one time, which may result in loose stools.

**CAUTIONS:** Mild, transient hypoglycemia may occur if taken on an empty stomach. Insulin dependent diabetics and pregnant women should consult their physician before use. Ribose may cause a transient increase in uric acid levels; therefore those that have chronic gout should consult their physician before use. Ribose may compete for phosphoglucomutase, a liver enzyme responsible for glycogen mobilization.

## **CONTRINDICATIONS**

Conditions of hypermagnesemia

## **STORAGE**

Store in a cool, dry place, away from direct light.  
Keep out of reach of children.

## REFERENCES

### BLOOD GLUCOSE

Eric R. Fenstad, Oladele Gazal, Linda M. Shecterle, J.A. St. Cyr & John G. Seifert: Dose Effects of D-Ribose on Glucose and Purine Metabolites: *The Internet Journal of Nutrition and Wellness*. 2008; Volume 5, Number 1.

Gross M, Zöllner N. *Klin Wochenschr*. 1991 Jan 4;69(1):31-6.

### SPORTS NUTRITION/SKELETAL MUSCLE

Hellsten Y, Skadhauge L, Bangsbo J. *Am J Physiol Regul Integr Comp Physiol*. 2004 Jan;286(1):R182-8.

D.Van Gammeran et al. *Current Therapeutic Research*. 2002. Vol 63.8.

Sahlin K, Tonkonogi M, Söderlund K. *Acta Physiol Scand*. 1998 Mar;162(3):261-6. Review.

Falk DJ, Heelan KA, Thyfault JP, Koch AJ. *J Strength Cond Res* 2003;17:810–16.

### FATIGUE SUPPORT

Teitelbaum JE, Johnson C, St Cyr J. *J Altern Complement Med*. 2006 Nov;12(9):857-62

Olsen NJ, Park JH. *Am J Med Sci*. 1998 Jun;315(6): Review

Russell IJ, Michalek JE, Flechas JD, Abraham GE. *J Rheumatol*. 1995 May;22(5):953-8.

†These statements have not been evaluated by the Food and Drug Administration.  
This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by  
Douglas Laboratories  
600 Boyce Road

Pittsburgh, PA 15205

800-245-4440

You trust Douglas Laboratories.



© 2011 Douglas Laboratories. All Rights Reserved